

DISCOMFORT/SWELLING

Aleve:

- Take 2 tablets 1 hour prior to your procedure with food (protein).
- Take 2 tablets within the hour after your procedure with 2 light glasses of water.
- Take 1 pill 3 times a day for 2 days to prevent any swelling or discomfort.

*After the 2 days, take as needed for discomfort

Anaprox:

- Take 1 tablet 2 hours prior to procedure with food (protein).
- Take 1 tablets 1 within the hour after your procedure with 2 large glasses of water.
- Take 1 pill 3 times a day for 2 days to prevent any swelling or discomfort.

*After the 2 days, take as needed for discomfort.

ANTIBIOTICS

Penicillin / Amoxicillin / Augmentin:

Take 2 tablets first dose In office At home
Then take 1 tablet every six hours (four times a day) until all are gone.

Erythromycin / Metronidazole (Flagyl):

Take 1 tablet to start with a full glass of water In office At home
Then take 1 tablet every eight hours (three times per day) until all are gone.

PERIDEX/PERIO GARD

- Oral rinse to be used twice a day at half strength to avoid staining the teeth.
- Gently keep in mouth for 1-2 minutes. Use until you return for follow up care.
- Swab with Q-tip or 308 end-tufted brush twice a day. Use until you return for follow up care.